



OxaGen 10 (Oxandrolone)

10 mg/each

Oral Tablets

FORMULA

Each uncoated tablet contains: Oxandrolone USP (10 mg), Excipients (q.s.)

DOSAGE AND USE

Therapy with anabolic steroids is adjunctive to and not a replacement for conventional therapy. The duration of therapy with Oxandrolone will depend on the response of the patient and the possible appearance of adverse reactions. Therapy should be intermittent.

The daily adult dosage is 2.5 mg to 20 mg given in 2 to 4 divided doses. The desired response may be achieved with as little as 2.5 mg or as much as 20 mg daily. A course of therapy of 2 to 4 weeks is usually adequate. This may be repeated intermittently as indicated.

INDICATIONS

OxaGen 10 is used in the treatment of catabolic processes or tissue reducers. Oxandrolone is indicated in conditions such as chronic infections, major surgery, (corticosteroid induced myopathy, decubitus ulcers, burns), or severe trauma, where it is required to stop the effects of catabolic processes or loss of proteins.

It is important to remember that this type of substances are adjuvants in these cases and do not replace conventional treatment of these disorders. It is also used for the treatment of Turner's Syndrome, to treat short stature observed in these patients.

MECHANISM OF ACTION

It reverses the catabolic processes and negative balance of nitrogen enhancing protein synthesis and stimulating the appetite if there is a simultaneous intake of calories and proteins.

CONTRAINDICATIONS

Oxandrolone is normally used for the treatment of symptoms associated with osteoporosis. However, the long-term use is not recommended.

ADVERSE EFFECTS

Virilization, acne, enlarged clitoris, hoarseness, menstrual irregularities, hair loss have been observed in women.

In prepubertal men, acne, penis enlargement, increased frequency of erections, abnormal hair growth have been observed.

In postpubertal men, bladder irritability (frequent urge to urinate), chest pain, breast enlargement, frequent erections have been observed. Less frequently cases of anemia, iron deficiency, loss of appetite, oedema, nausea, vomit, unusual bleeding have been observed.

Hypercalcemia (depression, unusual tiredness), unusual darkening of skin.

In elderly patients prostate carcinoma or prostate hypertrophy (difficulty or frequent urge to urinate) can be seen.

In rare cases hepatic necrosis (black stools, headache, vomiting blood, bad breath), hepatic cancer (stomach ache or abdominal pain, unexplained weight loss), hepatic peliosis (dark urine, loss of appetite, red spots on the skin, mouth, nose, throat, fever, nausea, vomits) have been observed.

DRUG INTERACTIONS

The use of anabolic steroids with adrenocorticosteroids, glucocorticosteroids, drugs or food that contain sodium can increase the possibility of oedema.

Concomitant use of oxandrolone with anticoagulants can increase the effect of the anticoagulant.

Anabolic steroids can decrease the concentration of glucose, diabetic patients should be monitored for probable symptoms of hypoglycemia and doses of hypoglycemic agents or insulin should be adjusted.

Concomitant use of anabolic steroids with hepatotoxic drugs can increase the hepatotoxicity. Patients with hepatic disorders should be carefully monitored.

PRECAUTIONS AND WARNINGS

Use with caution in children and adolescents because of an early closure of the epiphysis. Medication should be withdrawn if there are no results after 6 months of treatment.

OVERDOSE

In case of adverse effects withdraw use of the drug and immediately call your doctor or pharmacist.

PACKAGING

Box with 50 tablets of 10 mg

STORAGE CONDITIONS

Store in a cool dry place below 30°C.

Protect from light. Keep out of reach of children.

Sale only with Prescription.

Made in India

MyoGen™ By Gallene Sciences PVT. LTD.

Ahmedabad, Gujarat 380061, India

myogenlabs.com